
Bibliography

- Akaike, H. (1973). *Information theory and an extension of the maximum likelihood principle*. In Petrov, B. and Csaki, F., editors, *Second International Symposium on Information Theory*, pages 267–281. Akademiai Kiado, Budapest, Hungary.
- Bahadur, R. (1961). *A representation of the joint distribution of responses to n dichotomous items*. In Solomon, H., editor, *Studies in Item Analysis and Prediction*, Stanford Mathematical Studies in the Social Sciences, volume VI. Stanford University Press, Stanford, California, USA.
- Barnett, A., Koper, N., Dobson, A., Schmiegelow, F., and Manseau, M. (2010). *Using information criteria to select the correct variance-covariance structure for longitudinal data in ecology*. *Methods in Ecology and Evolution*, 1(1):15–24.
- Bellamy, S., Gibberd, R., Hancock, L., Howley, P., Kennedy, B., Klar, N., Lipsitz, S., and Ryan, L. (2000). *Analysis of dichotomous outcome data for community intervention studies*. *Statistical Methods in Medical Research*, 2:135–160.
- Bellamy, S., Li, Y., Lin, X., and Ryan, L. (2005). *Quantifying pql bias in estimating cluster-level covariate effects in group-randomized trials*. *Statistica Sinica*, 15(4):1015–1032.
- Bilker, W., Brensinger, C., and Gur, R. (2004). *A two factor anova-like test for correlated correlations: Coranova*. *Multivariate Behavioral Research*, 39(4):565–594.
- Breitung, J., Chaganty, N., Daniel, R., Kenward, M., Lechner, M., Martus, P., Sabo, R., Wang, Y., and Zorn, C. (2010). *Discussion of “generalized estimating equations: Notes on the choice of the working correlation matrix”*. *Methods of Information in Medicine*, 49(5):426–432.
- By, K., Qaqish, B., Pressier, J., Perin, J., and Zink, R. (2011). *Orth: R and sas software for regression models of correlated binary data based on orthogonalized residuals and alternating logistic regressions*. The University of North Carolina at Chapel Hill Department of Biostatistics Technical Report Series. Working Paper 22.
- Carey, V., Zeger, S., and Diggle, P. (1993). *Modelling multivariate binary data*

- with alternating logistic regressions. *Biometrika*, 80(3):517–526.
- Chaganty, C. and Mav, D. (2007). Estimation methods for analyzing longitudinal data occurring in biomedical research. In Khattree, R. and Naik, D., editors, *Computational Methods in Biomedical Research*, pages 371–400. Chapman and Hall/CRC Press, Boca Raton, Florida, USA.
- Chaganty, N. (1997). An alternative approach to the analysis of longitudinal data via generalized estimating equations. *Journal of Statistical Planning and Inference*, 63(1):39–54.
- Chaganty, N. and Deng, Y. (2007). Ranges of measures of association for familial binary variables. *Communications in Statistics- Theory and Methods*, 36(3):587–598.
- Chaganty, N. and Joe, H. (2004). Efficiency of generalized estimating equations for binary responses. *Journal of the Royal Statistical Society: Series B (Statistical Methodology)*, 66(4):851–860.
- Chaganty, N. and Joe, H. (2006). Range of correlation matrices for dependent bernoulli random variables. *Biometrika*, 93(1):197–206.
- Chaganty, N. and Naik, D. (2002). Analysis of multivariate longitudinal data using quasi-least squares. *Journal of Statistical Planning and Inference*, 103(1):421–436.
- Chaganty, N. and Shults, J. (1999). On eliminating the asymptotic bias in the quasi-least squares estimate of the correlation parameter. *Journal of Statistical Planning and Inference*, 76(1-2):145–161.
- Christensen, R. (1987). *Plane Answers to Complex Questions: the Theory of Linear Models*. Springer-Verlag Inc., New York, New York, USA.
- Christensen, R. (1997). *Linear Models for Multivariate, Time Series, and Spatial Data*. Springer-Verlag Inc., New York, New York, USA.
- Crowder, M. (1995). On the use of a working correlation matrix in using generalised linear models for repeated measures. *Biometrika*, 82(2):407–410.
- Cui, J. (2007). *Qic program and model selection in gee analyses*. *Stata Journal*, 7(2):209–220.
- Dahmen, G., Rochon, J., Konig, I. R., and Ziegler, A. (2004). Sample size calculations for controlled clinical trials using generalized estimating equations (gee). *Methods of Information in Medicine*, 43(5):451–461.
- Davis, C. (2002). *Statistical Methods for the Analysis of Repeated Measurements*. Springer-Verlag Inc., New York, New York, USA.
- De Backer, M., De Keyser, P., De Vroey, C., and Lesaffre, E. (1996). *British Journal of Dermatology*, 134:16–17.
- Diggle, P., Heagerty, P., Liang, K., and Zeger, S. (2002). *Analysis of Longitudinal Data*, Second Edition. Springer-Verlag Inc., New York, New York,

USA.

- Diggle, P., Liang, K., and Zeger, S. (1994). *Analysis of Longitudinal Data*, First Edition. Springer-Verlag Inc., New York, New York, USA.
- Dobson, A. and Barnett, A. (2008). *An Introduction to Generalized Linear Models*, Third Edition. Chapman and Hall/CRC Press, Boca Raton, Florida, USA.
- Dunlop, D. (1994). *Regression for longitudinal data: A bridge from least squares regression*. *The American Statistician*, 48(4):299–303.
- Ellenberg, J. H. (1994). *Selection bias in observational and experimental studies*. *Statistics in Medicine*, 13(5-7):557–567.
- Ellenberg, S. S. (2000). *Placebo-controlled trials and active control trials in the evaluation of new treatments*. *Annals of Internal Medicine*, 133:464–470.
- Feng, R., McClure, L. A., Tiwari, H. K., and Howard, G. (2009). *A new estimate of family disease history providing improved prediction of disease risks*. *Statistics in Medicine*, 28(8):1269–1283.
- Fitzmaurice, G. (1995). *A caveat concerning independence estimating equations with multivariate binary data*. *Biometrics*, 51(1):309–317.
- French, B. and Heagerty, P. J. (2008). *Analysis of longitudinal data to evaluate a policy change*. *Statistics in Medicine*, 27(24):5005–5025.
- Galecki, A. (1994). *General class of covariance structures for two or more repeated factors in longitudinal data analysis*. *Communications in Statistics*, 23:3105–20.
- Gantz, B., Tyler, R., Knutson, J., Woodworth, G., Abbas, P., McCabe, B., Hinrichs, J., Tye-Murray, N., Lansing, C., Kuk, F., and Brown, C. (1988). *Evaluation of five different cochlear implant designs: audiologic assessment and predictors of performance*. *Laryngoscope*, 98(10):1100–1106.
- Godambe, V. (2002). *Estimating Functions*. Oxford University Press, Oxford, UK.
- Grady, J. and Helms, R. (1995). *Model selection techniques for the covariance matrix for incomplete longitudinal data*. *Statistics in Medicine*, 13(14):1397–1416.
- Guerra, M. and Shults, J. (2013). *On the simulation of longitudinal discrete data with specified marginal means and first-order antedependence*. *The American Statistician* (tentatively accepted).
- Guerra, M. and Shults, J. (forthcoming 2014). *Logistic Regression for Longitudinal Data*. Springer, New York, New York, USA.
- Guerra, M., Shults, J., Amsterdam, J., and Ten-Have, T. (2012). *The analysis of binary longitudinal data with time-dependent covariates*. *Statistics in*

Medicine, 10:931–948.

- Guo, W., Ratcliffe, S. J., and TenHave, T. T. (2004). A random pattern-mixture model for longitudinal data with dropouts. *Journal of the American Statistical Association*, 99:929–937.
- Halekoh, U., Højsgaard, S., and Yan, J. (2006). The *r* package *geepack* for generalized estimating equations. *Journal of Statistical Software*, 15(2):1–11.
- Hammill, B. and Preisser, J. (2006). A *sas/iml* software program for *gee* and regression diagnostics. *Computational Statistics and Data Analysis*, 51(2):1197–1212.
- Hardin, J. and Hilbe, J. (2003). *Generalized Estimating Equations*. Chapman and Hall/CRC Press, Boca Raton, Florida, USA.
- Hardin, J. and Hilbe, J. (2007). *Generalized Linear Models and Extensions*, Second Edition. Stata Press, College Station, Texas, USA.
- Hardin, J. and Hilbe, J. (2012). *Generalized Estimating Equations*, Second Edition. Chapman and Hall/CRC Press, Boca Raton, Florida, USA.
- Hedeker, D. and Gibbons, R. (2006). Missing Data in Longitudinal Studies, pages 279–312. John Wiley and Sons, Inc.
- Heitjan, D. F. and Basu, S. (1996). Distinguishing “missing at random” and “missing completely at random”. *The American Statistician*, 50(3):pp. 207–213.
- Hilbe, J. (2009). *Logistic Regression Models*. Chapman and Hall/CRC Press, Boca Raton, Florida, USA.
- Hin, L. and Wang, Y. (2009). Working-correlation-structure identification in generalized estimating equations. *Statistics in Medicine*, 28:642–658.
- Horton, N. J., Bechuk, J. D., Jones, C. L., Lipsitz, S. R., Catalano, P. J., Zahner, G. E. P., and Fitzmaurice, G. M. (1999). Goodness-of-fit for *gee*: an example with mental health service utilization. *Statistics in Medicine*, 18(2):213–222.
- Huber, P. (1967). The behavior of maximum likelihood estimation under non-standard conditions. In *Proceedings of the Fifth Berkeley Symposium on Mathematical Statistics and Probability*, Berkely, California, USA, pages 221–233.
- Hurvich, C. and Tsai, C.-L. (1989). Regression and time series model selection in small samples. *Biometrika*, 76(2):297–307.
- Jung, S. and Ahn, C. (2003). Sample size estimation for *gee* method for comparing slopes in repeated measurements data. *Statistics in Medicine*, 22(8):130–1515.
- Kauermann, G. and Carroll, R. (2001). A note on the efficiency of sandwich

- covariance matrix estimation*. *Journal of American Statistical Association*, 96:1387–1396.
- Kim, H., Hilbe, J., and Shults, J. (2008a). *On the designation of the patterned associations for longitudinal bernoulli data: weight matrix versus true correlation structure?*. UPenn Biostatistics Working Papers. Working Paper 29, 26.
- Kim, H. and Shults, J. (2010). *% qls sas macro: A sas macro for analysis of longitudinal data using quasi-least squares*. *Journal of Statistical Software*, 35(2):1–22.
- Kim, H. and Shults, J. (2013). *Analysis of unbalanced simultaneously clustered and longitudinal data using quasi-least squares in sas*. Under Review.
- Kim, H., Shults, J., Patterson, S., and Goldberg-Alberts, R. (2008b). *Analysis of adverse events in drug safety: A multivariate approach using stratified quasi-least squares*. UPenn Biostatistics Working Papers. Working Paper 29.
- Kumanyika, S., Wadden, T., Shults, J., Fassbender, J., Brown, S., Bowman, M., Brake, V., West, W., Frazier, J., Whitt-Glover, M., Kallan, M., Desnoee, E., and Wu, X. (2009). *Trial of family and friend support for weight loss in african american adults*. *Archives of Internal Medicine*, 169(19):1795–1804.
- Lefkopoulou, M., Moore, D., and Ryan, L. (1989). *The analysis of multiple correlated binary outcomes: Application to rodent teratology experiments*. *Journal of the American Statistical Association*, 84:810–815.
- Liang, K. and Zeger, S. (1986). *Longitudinal data analysis using generalized linear models*. *Biometrika*, 73(1):13–22.
- Liang, K., Zeger, S., and Qaqish, B. (1992). *Multivariate regression analysis for categorical data*. *Journal of Royal Statistical Society, Series B*, 54(1):3–40.
- Liao, K., Freres, D. R., and Troxel, A. B. (2012). *A transition model for quality-of-life data with non-ignorable non-monotone missing data*. *Statistics in Medicine*, 31(28):3444–3466.
- Little, R. and Rubin, D. (2002). *Statistical Analysis with Missing Data* (Second Edition). *John Wiley and Sons, Inc.*
- Lu, B., Preisser, J., Qaqish, B., Suchindran, C., Bangdiwala, S., and Wolfson, M. (2007). *A comparison of two bias-corrected covariance estimators for generalized estimating equations*. *Biometrics*, 63(3):935–941.
- Lu, N. and Zimmerman, D. (2005). *The likelihood ratio test for a separable covariance matrix*. *Statistics and Probability Letters*, 73:449–457.
- Magnus, P., Gjessing, H., Skrondal, A., and Skjaerven, R. (2001). *Paternal contribution to birth weight*. *Journal of Epidemiology and Community Health*,

55:873–877.

- Mancl, L. and DeRouen, T. (2001). *A covariance estimator for gee with improved small-sample properties*. *Biometrics*, 57(1):126–134.
- McCullagh, P. and Nelder, J. (1989). *Generalized Linear Models*, Second Edition. *Chapman and Hall/CRC Press Monographs on Statistics and Applied Probability*, Boca Raton, Florida, USA.
- Miller, M., Davis, C., and Landis, J. (1993). *The analysis of longitudinal polytomous data: generalized estimating equations and connections with weighted least squares*. *Biometrics*, 49(4):1033–1044.
- Molenberghs, G. (2010). *Editorial. “generalized estimating equations: Notes on the choice of the working correlation matrix”*. *Methods of Information in Medicine*, 49(5):419–420.
- Molenberghs, G. and Kenward, M. (2010). *Semi-parametric marginal models for hierarchical data and their corresponding full models*. *Computational Statistics & Data Analysis*, 54(2):585–597.
- Molenberghs, G. and Verbeke, G. (2005). *Models for Discrete Longitudinal Data*. *Springer, New York, New York, USA*.
- Monahan, J. (2008). *A Primer on Linear Models*. *Chapman and Hall/CRC Press, Boca Raton, Florida, USA*.
- Naik, D. and Rao, S. (2001). *Analysis of multivariate repeated measures data with a kronecker product structured covariance matrix*. *Journal of Applied Statistics*, 28:91–105.
- Nelder, J. and Wedderburn, R. (1972). *Generalized linear models*. *Journal of the Royal Statistical Society, Series A*, 135(3):370–384.
- Newton, H. (1988). *TIMESLAB: A Time Series Analysis Laboratory*. *Wadsworth and Brooks/Cole, Pacific Grove, California, USA*.
- Nunez-Anton, V. and Woodworth, G. (1994). *Analysis of longitudinal data with unequally spaced observations and time-dependent correlated errors*. *Biometrics*, 50(2):445–456.
- O’Brien, S., Holubkov, R., and Reis, E. (2004). *Identification, evaluation, and management of obesity in an academic primary care*. *Pediatrics*, 114(e):154–159.
- Olkin, I. and Pratt, J. W. (1958). *A multivariate tchebycheff inequality*. *The Annals of Mathematical Statistics*, 29(1):226–234.
- Pan, W. (2001). *Akaike’s information criterion in generalized estimating equations*. *Biometrics*, 57:120–125.
- Park, T. (1993). *A comparison of the generalized estimating equation approach with the maximum likelihood approach for repeated measurements*. *Statistics in Medicine*, 12(18):1723–1732.

- Preisser, J., Arcury, T., and Quandt, S. (2003a). *Detecting patterns of occupational illness clustering with alternating logistic regressions applied to longitudinal data*. *American Journal of Epidemiology*, 158(5):495–501.
- Preisser, J., Qaqish, B., and Perin, J. (2008). *A note on deletion diagnostics for estimating equations*. *Biometrika*, 95(2):509–513.
- Preisser, J. and Qaqish, B. F. (1996). *Deletion diagnostics for generalised estimating equations*. *Biometrika*, 83(3):551–562.
- Preisser, J., Reboussin, B., Song, E., and Wolfson, M. (2007). *The importance and role of intracluster correlations in planning cluster trials*. *Epidemiology*, 18(5):552–560.
- Preisser, J., Young, M., Zaccaro, D., and Wolfson, M. (2003b). *An integrated population-averaged approach to the design, analysis, and sample size determination of cluster-unit trials*. *Statistics in Medicine*, 22(8):1235–1254.
- Preisser, J. S., By, K., Perin, J., and Qaqish, B. F. (2012). *Deletion diagnostics for alternating logistic regressions*. *Biometrical Journal*, 54(5):701–715.
- Preisser, J. S., Lohman, K., and Rathouz, P. (2002). *Longitudinal studies with dropouts missing at random*. *Statistics in Medicine*, 21:3035–3054.
- Prentice, R. (1988). *Correlated binary regression with covariates specific to each binary observation*. *Biometrics*, 44(4):1033–1048.
- Putt, M. E. (2006). *Power to detect clinically relevant carry-over in a series of cross-over studies*. *Statistics in Medicine*, 25(15):2567–2586.
- Qaqish, B. (2003). *A family of multivariate binary distributions for simulating correlated binary variables with specified marginal means and correlations*. *Biometrika*, 90:455–463.
- Qaqish, B. and Liang, K. (1992). *Marginal models for correlated binary responses with multiple classes and multiple levels of nesting*. *Biometrics*, 48(3):939–950.
- Qaqish, B., Zink, R., and Preisser, J. (2012). *Orthogonalized residuals for estimation of marginally specified association parameters in multivariate binary data*. *Scandinavian Journal of Statistics*, 39(3):515–527.
- Qu, A. and Li, R. (2006). *Quadratic inference functions for varying coefficient models with longitudinal data*. *Biometrics*, 62(2):379–391.
- Qu, A. and Lindsay, B. (2003). *Building adaptive estimating equations when inverse of covariance estimation is difficult*. *Journal of the Royal Statistical Society, Series B*, 65(1):127–142.
- Qu, A., Lindsay, B., and Li, B. (2000). *Improving generalised estimating equations using quadratic inference functions*. *Biometrika*, 87(4):823–836.
- Qu, A. and Song, P. (2004). *Assessing robustness of generalised estimating equations and quadratic inference functions*. *Biometrika*, 91(2):447–459.

- Qu, Y., Williams, G., Beck, G., and Medendorp, S. (1992). *Latent variable models for clustered dichotomous data with multiple subclusters*. *Biometrics*, (4):1095–1102.
- Ratcliffe, S. and Shults, J. (2008). *Geeqbox: A matlab toolbox for implementation of quasi-least squares and generalized estimating equations*. *Journal of Statistical Software*, 25(14):1–13.
- Rencher, A. and Schaalje, G. (2007). *Linear Models in Statistics, Second Edition*. Wiley-Interscience, Hoboken, New Jersey, USA.
- Rochon, J. (1998). *Application of gee procedures for sample size calculations in repeated measures experiments*. *Statistics in Medicine*, 17(14):1643–1658.
- Rotnitzky, A. and Jewell, N. (1990). *Hypothesis testing of regression parameters in semiparametric generalized linear models for cluster correlated data*. *Biometrika*, 77:485–497.
- Roy, A., Bhaumik, D., Subhash, S., and Gibbons, R. (2007). *Sample size determination for hierarchical longitudinal designs with differential attrition rates*. *Biometrics*, 63(3):699–707.
- Roy, A. and Khattree, R. (2003). *Tests for mean and covariance structures relevant in repeated measures based discriminant analysis*. *Journal of Applied Statistical Science*, 12(2):91–104.
- Roy, A. and Khattree, R. (2005). *On implementation of a test for kronecker product covariance structure for multivariate repeated measures data*. *Statistical Methodology*, 2:297–306.
- Rubin, D. B. (1976). *Inference and missing data*. *Biometrika*, 63(3):581–592.
- Sabo, R. and Chaganty, N. (2009). *Adaptation of quasi-least squares to estimate correlations within a nuclear family*. *Communications in Statistics: Theory and Methods*, 38(16):3059–3076.
- Sabo, R. and Chaganty, N. (2010a). *Estimation methods for an autoregressive familial correlation structure*. *Communications in Statistics: Theory and Methods*, 39(6):973–991.
- Sabo, R. and Chaganty, N. (2010b). *Hypothesis testing for various familial dependence structures*. *Communications in Statistics: Simulation and Computation*, 13(1):207–219.
- Shaw, P. A. and Proschan, M. A. (2013). *Null but not void: considerations for hypothesis testing*. *Statistics in Medicine*, 32(2):196–205.
- Shi, G. and Chaganty, N. (2004). *Application of quasi-least squares to analyze replicated autoregressive time series regression models*. *Journal of Applied Statistics*, 31(10):1147–1156.
- Shih, W. (1997). *Sample size and power calculation for periodontal and other*

- studies with clustered samples using the method of generalized estimating equations.* *Biometrical Journal*, 39(8):899–908.
- Shults, J. (1996). The Analysis of Unbalanced and Unequally Spaced Longitudinal Data using Quasi-Least Squares. *PhD thesis, Department of Mathematics and Statistics, Old Dominion University, Norfolk, Virginia.*
- Shults, J. (2011). *Generalized estimating equations: Notes on the choice of the working correlation matrix - continued. letter to the editor.* *Methods of Information in Medicine*, 50(1):96–99.
- Shults, J. and Chaganty, N. (1998). *Analysis of serially correlated data using quasi-least squares.* *Biometrics*, 54(4):1622–1630.
- Shults, J., Mazurick, C., and Landis, J. (2006a). *Analysis of repeated bouts of measurements in the framework of generalized estimating equations.* *Statistics in Medicine*, 25(23):4114–4128.
- Shults, J. and Morrow, A. (2002). *Use of quasi-least squares to adjust for two levels of correlation.* *Biometrics*, 58(3):521–530.
- Shults, J. and Ratcliffe, S. (2009). *Analysis of multi-level correlated data in the framework of generalized estimating equations via xtmultcorr procedures in stata and qls functions in matlab.* *Statistics and Its Interface*, 2:187–196.
- Shults, J., Ratcliffe, S., and Leonard, M. (2007). *Improved generalized estimating equation analysis via xtqls for implementation of quasi-least squares in stata.* *Stata Journal*, 7(2):147–166.
- Shults, J., Sun, W., Tu, X., Kim, H., Amsterdam, J., Hilbe, J., and Ten-Have, T. (2009). *A comparison of several approaches for choosing between working correlation structures in generalized estimating equation analysis of longitudinal binary data.* *Statistics in Medicine*, 28(8):2338–2355.
- Shults, J., Tu, X., and Amsterdam, J. (2006b). *On the violation of bounds for the correlation in generalized estimating equation analyses of binary data from longitudinal trials.* UPenn Biostatistics Working Papers. Working Paper 8.
- Shults, J., Whitt, M., and Kumanyika, S. (2004). *Analysis of data with multiple sources of correlation in the framework of generalized estimating equations.* *Statistics in Medicine*, 23(20):3209–3226.
- Song, P. (2007). *Correlated Data Analysis: Modeling, Analytics, and Applications.* *Springer Advances in Statistics, New York, New York, USA.*
- Spiegelhalter, D. J., Best, N. G., Carlin, B. P., and Van Der Linde, A. (2002). *Bayesian measures of model complexity and fit.* *Journal of the Royal Statistical Society: Series B (Statistical Methodology)*, 64(4):583–639.
- Stokes, M., Davis, C., and Koch, G. (2000). *Categorical Data Analysis Using the SAS System*, 2nd edition. Cary, North Carolina: SAS Institute Inc.
- Sugiura, N. (1978). *Further analysis of the data by akaike's information cri-*

- terion and the finite corrections.* Communications in Statistics, Theory and Methods, *A(7)*:13–26.
- Sun, W., Shults, J., and Leonard, M. (2009). *A note on the use of unbiased estimating equations to estimate correlation in analysis of longitudinal trials.* Biometrical Journal, *51(1)*:5–18.
- Sutradhar, B. and Das, K. (1999). *On the efficiency of regression estimators in generalised linear models for longitudinal data.* Biometrika, *86(2)*:459–465.
- Sutradhar, B. and Das, K. (2000). *On the accuracy of efficiency of estimating equation approach.* Biometrics, *56(2)*:622–625.
- Teerenstra, S., Lu, B., Preisser, J., Achterberg, T., and Borm, G. (2010). *On the accuracy of efficiency of estimating equation approach.* Biometrics, *66(10)*:1230–1237.
- Troxel, A. B., Fairclough, D. L., Curran, D., and Hahn, E. A. (1998). *Statistical analysis of quality of life with missing data in cancer clinical trials.* Statistics in Medicine, *17(5-7)*:653–666.
- Troxel, A. B., Lipsitz, S. R., Fitzmaurice, G. M., Ibrahim, J. G., Sinha, D., and Molenberghs, G. (2010). *A weighted combination of pseudo-likelihood estimators for longitudinal binary data subject to non-ignorable non-monotone missingness.* Statistics in Medicine, *29(14)*:1511–1521.
- Tu, X., Kowalski, J., Zhang, J., K.G., L., and P., C.-C. (2004). *Power analyses for longitudinal trials and other clustered designs.* Statistics in Medicine, *23(18)*:2799–2815.
- Tu, X., Zhang, J., Kowalski, J., Shults, J., Feng, C., Sun, W., and Tang, W. (2007). *Power analyses for longitudinal study designs with missing data.* Statistics in Medicine, *26(15)*:2958–2981.
- Wang, Y. and Carey, V. (2003). *Working correlation structure misspecification, estimation and covariate design: Implications for generalised estimating equations performance.* Biometrika, *90(1)*:29–41.
- Wedderburn, R. (1974). *Quasi-likelihood functions, generalized linear models, and the gaussian method.* Biometrika, *61(3)*:439–447.
- Weissfeld, L. and Kshirsagar, A. (1992). *A modified growth curve model and its application to clinical studies.* Australian and New Zealand Journal of Statistics, *34(2)*:161–168.
- Westgate, P. M. (2013). *A bias correction for covariance estimators to improve inference with generalized estimating equations that use an unstructured correlation matrix.* Statistics in Medicine, *32(16)*:2850–2858.
- White, H. (1982). *Maximum likelihood estimation of misspecified models.* Econometrica, *50(1)*:1–25.

- White, M. (2011). *Determinants of change in vitamin d level in chronic kidney disease*. Master's thesis, University of Pennsylvania, Philadelphia, Pennsylvania.
- Whittle, P. (1958). *A multivariate generalization of tchevichev's inequality*. The Quarterly Journal of Mathematics, 9(1):232–240.
- Xie, J. and Shults, J. (2009). *Implementation of quasi-least squares with the r package qlspack*. UPenn Biostatistics Working Papers. Working Paper 32.
- Xie, J., Shults, J., Peet, J., Stambolian, D., and Cotch, M. (2010). *Quasi-least squares with mixed linear correlation structures*. Statistics and Its Interface, 3(2):223–234.
- Yan, J. (2002). *geepack: yet another package for generalized estimating equations*. R News, 2:12–14.
- Zeger, S., Liang, K., and Albert, P. (1988). *Models for longitudinal data: a generalized estimating equation approach*. Biometrics, pages 1049–60.
- Zeger, S., Liang, K., and Self, S. (1985). *The analysis of binary longitudinal data with time-independent covariates*. Biometrika, 72(3):31–38.
- Zhao, L. P. and Prentice, R. (1990). *Longitudinal data analysis for discrete and continuous outcomes*. Biometrika, 77(3):642–648.
- Zheng, B. (2000). *Summarizing the goodness of fit of generalized linear models for longitudinal data*. Statistics in Medicine, 19:1265–1275.
- Ziegler, A. (2011). *Generalized Estimating Equations*. Springer, New York, Heidelberg, Dordrecht, London.
- Ziegler, A. and Vens, M. (2010). *Generalized estimating equations: Notes on the choice of the working correlation matrix*. Methods of Information in Medicine, 49(5):421–425.